



Pronounced ev-RIZ-dee.

Learn more about Evrysdi

Proven to make a difference in infants, children, and adults with SMA

In people 2 months and older who have a broad
range of symptoms and motor function

What is Evrysdi?

Evrysdi is a prescription medicine used to treat spinal muscular atrophy (SMA) in adults and children 2 months of age and older. It is not known if Evrysdi is safe and effective in children under 2 months of age.

Important Safety Information

- Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:
 - have liver problems

Please see additional Important Safety Information and full Prescribing Information.



About
Evrysdi

How Evrysdi
was studied

Results in
Type 1 SMA

Results in
Type 2 & 3
SMA

Safety
studied

How to take
Evrysdi

MySMA
Support

Talking to
your doctor

Summary

Important
Safety
Information

Prescribing
Information

Get to know **Evrysdi**

This brochure was developed to provide you with information about Evrysdi, including:



How Evrysdi was studied



Results in infants (Type 1 SMA)



Results in children and adults (Type 2 and 3 SMA)



Safety studied



How to take Evrysdi



MySMA Support services



Questions to ask your healthcare provider



Although this brochure is meant to provide helpful information, it does not replace conversations with your healthcare provider.

Please see additional Important Safety Information and full Prescribing Information.



**About
Evrysdi**

How Evrysdi
was studied

Results in
Type 1 SMA

Results in
Type 2 & 3
SMA

Safety
studied

How to take
Evrysdi

MySMA
Support

Talking to
your doctor


Summary

Important
Safety
Information

Prescribing
Information


Evrysdi is a disease-modifying therapy **approved to treat SMA**

SMA is a genetic condition caused by the shortage of a key protein called SMN, or survival motor neuron, which is needed for nerves and muscles to function properly.



Evrysdi is designed to work by helping to make and maintain more SMN protein

In animal studies, Evrysdi was distributed throughout the body



Evrysdi increased and maintained levels of SMN protein

Within 4 weeks of starting Evrysdi, median* SMN protein levels

MORE THAN DOUBLED

and

WERE MAINTAINED

throughout 12 months of studies across all SMA types

*A median is the middle value of a set of data that have been put into numerical order. The median is the value that divides the data into two halves.

Important Safety Information

- Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:
 - are pregnant or plan to become pregnant. If you are pregnant, or are planning to become pregnant, ask your healthcare provider for advice before taking this medicine. Evrysdi may harm your unborn baby.

Please see additional Important Safety Information and full Prescribing Information.



**About
Evrysdi**

How Evrysdi
was studied

Results in
Type 1 SMA

Results in
Type 2 & 3
SMA

Safety
studied

How to take
Evrysdi

MySMA
Support

Talking to
your doctor


Summary

Important
Safety
Information

Prescribing
Information


Evrysdi is being studied in a broad range of infants, children, and adults with SMA

Including more than 450 people with Type 1, 2, or 3 SMA from:



Efficacy was only studied in people from 2 months to 25 years old


with the following characteristics:



Those with varying symptoms and motor functions*



Those who could or could not walk



Those without scoliosis to those with severe scoliosis or joint contractures



Those who have or have not taken medication for SMA before

This is the first time a medication for SMA is being studied in people over 18 years of age, some of whom have severe scoliosis

*Motor function was measured using several tools including the Bayley Scales of Infant and Toddler Development Gross Motor Scale-Third Edition (BSID-III), the Motor Function Measure-32 Items (MFM-32), and the Revised Upper Limb Module (RULM) assessment.

Important Safety Information

- Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:
 - are a woman who can become pregnant:
 - Before you start your treatment with Evrysdi, your healthcare provider may test you for pregnancy. Because Evrysdi may harm your unborn baby, your healthcare provider will decide if taking Evrysdi is right for you during this time
 - Talk to your healthcare provider about birth control methods that may be right for you. Use birth control while on treatment and for at least 1 month after stopping Evrysdi

Please see additional Important Safety Information and full Prescribing Information.



About
Evrysdi

**How Evrysdi
was studied**

Results in
Type 1 SMA

Results in
Type 2 & 3
SMA

Safety
studied

How to take
Evrysdi

MySMA
Support

Talking to
your doctor

Summary

Important
Safety
Information

Prescribing
Information

Evrysdi received FDA approval based on results from the first 12 months of **2 main ongoing studies**



FIREFISH (infants with Type 1 SMA)

FIREFISH is an open-label study made up of 2 parts

- 62 infants (2 to 7 months old) with Type 1 SMA were studied
- The main measurement was the ability of infants to sit without support for at least 5 seconds* and survival without permanent breathing support† after 12 months of taking Evrysdi
 - Part 1 includes 21 infants. It explored the recommended dose of Evrysdi for people in this age group. The study also measured the safety and effectiveness of Evrysdi after 12 months. The results from Part 1 can be seen on pages 6 and 7 of this brochure
 - Part 2 includes 41 infants and is measuring the safety and effectiveness of Evrysdi at the recommended dose



SUNFISH (children and adults with Type 2 or 3 SMA)

SUNFISH is a randomized, placebo-controlled study made up of 2 parts

- 231 children and adults (2 to 25 years old) with Type 2 or 3 SMA and a range of motor functions were studied:
- The main measurement was the change in motor function after 12 months of treatment,‡ compared with those not taking Evrysdi§
 - Part 1 includes 51 children and adults and explored the recommended dose. It is also exploring the safety of Evrysdi for people in this age group. This part includes 7 people who could walk
 - Part 2 includes 180 people and is measuring the safety and effectiveness of Evrysdi at the recommended dose (n=120), compared with those not taking Evrysdi (n=60). These results can be found on pages 8-11 of this brochure. Part 2 includes:
 - Those with scoliosis (n=120), sometimes severe (n=57)
 - Those with or without contractures
 - Those who cannot walk (n=180)

A third, supportive study is currently looking at the safety of Evrysdi in people previously treated for SMA



JEWELFISH: A study to monitor safety for people with Type 1, 2, or 3 SMA who were previously treated with other SMA medications

This study includes 174 people from 1 to 60 years old who had taken other SMA medications before. This study is ongoing.

*Measured by Item 22 of the BSID-III Gross Motor Scale.

†Permanent support was defined as having a tracheostomy (a surgery where a tube is inserted in the front of the throat into the windpipe) or more than 21 days of noninvasive ventilation support (16 or more hours a day) or being intubated (a procedure where a breathing tube is inserted down the throat and into the windpipe) to help with breathing in the absence of an acute reversible event.

‡Measured by the MFM-32 scale.

§Those not taking Evrysdi were on placebo, a substance that has no active medication and is often used in studies.

FDA stands for the US Food and Drug Administration.

Please see additional Important Safety Information and full Prescribing Information.



About
Evrysdi

**How Evrysdi
was studied**

Results in
Type 1 SMA

Results in
Type 2 & 3
SMA

Safety
studied

How to take
Evrysdi

MySMA
Support

Talking to
your doctor

Summary

Important
Safety
Information

Prescribing
Information

Evrysdi helped infants **achieve a key motor milestone**

Out of the infants taking the recommended dose of Evrysdi for 12 months



Untreated infants with Type 1 SMA are not expected to sit without support

- The infants in this part of the study ranged in age from 3 to 7 months at the start of treatment, but many were closer to 7 months



How were these results measured?

The results shown here were measured using a tool called the Bayley Scales of Infant and Toddler Development Gross Motor Scale-Third Edition (BSID-III), which is used to measure sitting (Item 22) and other developmental movements.

Important Safety Information

- Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:
 - are an adult male planning to have children: Evrysdi may affect a man's ability to have children (fertility). If this is of concern to you, make sure to ask a healthcare provider for advice

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



About
Evrysdi

How Evrysdi
was studied

**Results in
Type 1 SMA**

Results in
Type 2 & 3
SMA

Safety
studied

How to take
Evrysdi

MySMA
Support

Talking to
your doctor

Summary

Important
Safety
Information

Prescribing
Information

Evrysdi helped infants survive without permanent breathing support

Out of the infants taking Evrysdi



Additional exploratory observations in eating and swallowing

This information is considered **exploratory**. This means it was not designed to show a treatment effect so conclusions cannot be drawn from these observational data.

Out of the infants taking the recommended dose of Evrysdi for 12 months:

- 88% were able to eat by mouth (15 out of 17 infants)[†]
- 88% were able to swallow (15 out of 17 infants)[‡]

*Permanent support was defined as having a tracheostomy (a surgery where a tube is inserted in the front of the throat into the windpipe) or more than 21 days of noninvasive ventilation support (16 or more hours a day) or being intubated (a procedure where a breathing tube is inserted down the throat and into the windpipe) to help with breathing in the absence of an acute reversible event.

[†]Two of these infants could eat some food by mouth, but also used a feeding tube.

[‡]One infant could not swallow at the start of the study.

Important Safety Information

- Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:
 - are breastfeeding or plan to breastfeed. It is not known if Evrysdi passes into breast milk and may harm your baby. If you plan to breastfeed, discuss with your healthcare provider about the best way to feed your baby while on treatment with Evrysdi

Please see additional Important Safety Information and full Prescribing Information.



About
Evrysdi

How Evrysdi
was studied

**Results in
Type 1 SMA**

Results in
Type 2 & 3
SMA

Safety
studied

How to take
Evrysdi

MySMA
Support

Talking to
your doctor

Summary

Important
Safety
Information


Prescribing
Information

After taking Evrysdi, **motor function improved in children and adults**

As measured by MFM-32

Children and adults treated with Evrysdi showed improved motor function compared with those not taking Evrysdi*

Change in motor function measured over 12 months



Talk to your healthcare provider about setting treatment goals specific to you. Goals may vary based on factors like your age and current motor function abilities.

*Those not taking Evrysdi were on placebo, a substance that has no active medication and is often used in studies.

†In some studies, including this one, if someone's data cannot be collected on time for any reason, their progress cannot be counted in that part of the study. This chart only includes the information that was collected on time.

MFM-32 stands for the Motor Function Measure-32 Items

Important Safety Information

- **Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine

Please see additional Important Safety Information and full Prescribing Information.



About
Evrysdi

How Evrysdi
was studied

Results in
Type 1 SMA

**Results in
Type 2 & 3
SMA**

Safety
studied

How to take
Evrysdi

MySMA
Support

Talking to
your doctor

Summary

Important
Safety
Information


Prescribing
Information

MFM-32 measures motor function abilities that relate to **important daily functions**

The MFM-32 scale includes motor movement across 3 categories:



Standing/transfer movements



Upper/lower body movements



Hand/foot movements



How were these results measured?

The results shown here were measured using a tool called the Motor Function Measure-32 items (MFM-32) scale, which is designed to capture change for a broad range of people, including those who can and cannot walk. It evaluates 32 different elements to assess head, trunk, and upper limb motor movement. The elements are grouped into 3 categories of movements, including standing/transfers, upper/lower body, and hand/foot.

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



About
Evrysdi

How Evrysdi
was studied

Results in
Type 1 SMA

**Results in
Type 2 & 3
SMA**

Safety
studied

How to take
Evrysdi

MySMA
Support

Talking to
your doctor

Summary


Important
Safety
Information

Prescribing
Information

After taking Evrysdi, **upper limb function improved in children and adults**

As measured by the RULM assessment

Children and adults treated with Evrysdi showed greater improvements in upper limb function compared with those who were not taking Evrysdi*



*Those not taking Evrysdi were on placebo, a substance that has no active medication and is often used in studies.

†In some studies, including this one, if someone's data cannot be collected on time for any reason, their progress cannot be counted in that part of the study. This chart only includes the information that was collected on time.

RULM stands for the Revised Upper Limb Module (RULM) assessment

Important Safety Information

- You should receive Evrysdi from the pharmacy as a liquid that can be given by mouth or through a feeding tube. The liquid solution is prepared by your pharmacist. If the medicine in the bottle is a powder, **do not use it**. Contact your pharmacist for a replacement

Please see additional Important Safety Information and full Prescribing Information.



About
Evrysdi

How Evrysdi
was studied

Results in
Type 1 SMA

**Results in
Type 2 & 3
SMA**

Safety
studied

How to take
Evrysdi

MySMA
Support

Talking to
your doctor

Summary

Important
Safety
Information

Prescribing
Information

RULM measures upper limb strength by **evaluating the performance of daily tasks**

The RULM assessment includes tests such as:




Picking up
coins/small objects



Using a
pencil



Raising a cup
to mouth



Opening a
plastic container



How were these results measured?

The results shown here were measured using a tool called the Revised Upper Limb Module (RULM) assessment. It evaluates strength in arm movements and the ability to perform tasks in people 2.5 years or older who have varying levels of muscle weakness. This measurement was specifically designed to test upper limb function in people living with SMA.

The RULM assessment also tests the ability to perform other activities, like:

- Bringing hands from lap to table
- Placing a coin or other small object into a cup
- Reaching to the side
- Bringing a hand to shoulder
- Pushing a button
- Tearing paper
- Lifting and moving ~1/2-lb and ~1-lb weights*
- Bringing ~1-lb weight* from lap to table or eye level
- Bringing both arms above head
- Bringing ~1-lb and ~2-lb weights* above shoulder
- Bringing hand above shoulder

*Weight is approximate and has been converted from grams to pounds.

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



About
Evrysdi

How Evrysdi
was studied

Results in
Type 1 SMA

**Results in
Type 2 & 3
SMA**

Safety
studied

How to take
Evrysdi

MySMA
Support

Talking to
your doctor

Summary

Important
Safety
Information

Prescribing
Information

Safety studied in infants, children, and adults

Evrysdi is being studied in people from 2 months to 60 years old with Type 1, 2, or 3 SMA

The most common side effects of Evrysdi include:

- **For infants, Type 1 SMA (infantile-onset):**

- fever
- diarrhea
- rash
- runny nose, sneezing, sore throat, and cough (upper respiratory infection)
- lung infection
- constipation
- vomiting

- **For children and adults, Type 2 or 3 SMA (later-onset):**

- fever
- diarrhea
- rash

These are not all of the possible side effects of Evrysdi. For more information, ask your healthcare provider.

No one stopped taking Evrysdi because of side effects in the main clinical studies after 12 months of taking Evrysdi.

Safety of Evrysdi in people previously treated with other SMA medications is currently being studied

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



About
Evrysdi

How Evrysdi
was studied

Results in
Type 1 SMA

Results in
Type 2 & 3
SMA

**Safety
studied**

How to take
Evrysdi

MySMA
Support

Talking to
your doctor

Summary

Important
Safety
Information

Prescribing
Information

The first and only medication **to treat SMA with at-home dosing**

Evrysdi is a liquid treatment, taken by mouth or feeding tube, that is:




Delivered directly to you



Stored in the refrigerator



Given after measuring your dose with the oral syringes provided



Taken once a day by mouth or feeding tube

Your healthcare provider will determine the right dose for you based on age and weight.
Always take the dose exactly as prescribed.

You can administer Evrysdi to yourself or with the help of the person who cares for you.
It should be taken around the same time each day after a meal.

Talk to your healthcare provider to learn more about Evrysdi

Please read the full [Patient Information](#) and [Instructions for Use](#) that come with your prescription before taking the first dose.

Important Safety Information

- Avoid getting Evrysdi on your skin or in your eyes. If Evrysdi gets on your skin, wash the area with soap and water. If Evrysdi gets in your eyes, rinse your eyes with water

Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



Your shipment is **delivered directly to your door** and will contain a 1-month supply of Evrysdi



Each package within the shipment includes:

- Up to 3 bottles of liquid medication with bottle adapter and cap
- Reusable oral syringes
- Instructions for Use
- Prescribing Information and Patient Information



Accredo* is the only specialty pharmacy that prepares and ships Evrysdi directly to you each month

Evrysdi is not available at retail pharmacies. Accredo will call you to coordinate each delivery on a day that works for you. Although Accredo is not part of Genentech, it is an important part of the MySMA Support team.

*Accredo Health Group, Inc., is an Express Scripts specialty pharmacy. Accredo is not part of Genentech and maintains independence in its operations as a specialty pharmacy and in its role as a healthcare provider.

Please see additional Important Safety Information and full Prescribing Information.



About
Evrysdi

How Evrysdi
was studied

Results in
Type 1 SMA

Results in
Type 2 & 3
SMA

Safety
studied

**How to take
Evrysdi**

MySMA
Support

Talking to
your doctor

Summary

Important
Safety
Information

Prescribing
Information

MySMA Support is **here for you**

Introducing a support service from Genentech that can help provide information about Evrysdi, including:



Product Education



Insurance and Financial Assistance



Additional Support



Coordination With Specialty Pharmacy for Delivery

MySMA Support, including the PAL, does not provide medical advice and is not a substitute for your medical team.
Your healthcare provider should always be your main resource for any questions about your health and medical care.

Please see additional Important Safety Information and full Prescribing Information.



Partnership and Access Liaisons (PALs) are here to help

A PAL is the local, main point of contact from Genentech who supports people living with SMA.
A PAL can:



Provide in-person or virtual support based on your preference



Explain insurance coverage and potential financial support options



Answer questions on Evrysdi, including how it works, how to take it, what results were seen in clinical trials, and important safety information



Refer you to helpful resources



Talk to a PAL in person, over the phone, or online. Connect with a local PAL by visiting www.Evrysdi.com/PAL or calling (833) 387-9734 (Monday-Friday, 9 AM to 8 PM ET)



Please see additional [Important Safety Information](#) and full [Prescribing Information](#).



About Evrysdi

How Evrysdi was studied

Results in Type 1 SMA

Results in Type 2 & 3 SMA

Safety studied

How to take Evrysdi

MySMA Support

Talking to your doctor

Summary

Important Safety Information

Prescribing Information

Talking to your healthcare provider about Evrysdi

If you're considering treatment for SMA, it may be the right time to talk with your healthcare provider about Evrysdi

In addition to your own questions, you can ask the following questions during your discussion:

About Evrysdi

- ☐ Was Evrysdi studied in people like me/my child?
- ☐ What kinds of results have people seen while taking Evrysdi?
- ☐ What is the important safety information that I should know about?
- ☐ How does Evrysdi work?
- ☐ Is Evrysdi right for me/my child?



Taking or Giving Evrysdi



- ☐ Where do I get Evrysdi?
- ☐ When do I take Evrysdi?
- ☐ How do I take or give Evrysdi?

Insurance Coverage and Support

- ☐ Where can I find more information about insurance coverage and financial support?
- ☐ Where can I find additional resources?

Bring this list to your next appointment with your healthcare provider, and visit www.Evrysdi.com for more information

Please see additional Important Safety Information and full Prescribing Information.



About
Evrysdi

How Evrysdi
was studied

Results in
Type 1 SMA

Results in
Type 2 & 3
SMA

Safety
studied

How to take
Evrysdi

MySMA
Support

**Talking to
your doctor**

Summary

Important
Safety
Information

Prescribing
Information

Proven to make a difference

in infants, children, and adults with SMA

Being studied in more than 450 people living with SMA



Evrysdi improved motor function scores in 2 main studies of infants, children, and adults:

- 41% of infants were able to sit for at least 5 seconds without support after 12 months (7 out of 17 infants)
- 90% of infants (19 out of 21) at 12 months and 81% of infants (17 out of 21) at 23 months were alive and could breathe without permanent support
- Motor function improved in children and adults after 12 months (average 1.36-point increase on the MFM-32 scale for those treated with Evrysdi vs average 0.19-point decrease for those not taking Evrysdi)



Evrysdi increased and maintained SMN protein levels in clinical studies

- Within 4 weeks, median SMN protein levels more than doubled and were maintained throughout the 12 months of studies



The safety of Evrysdi is being studied in people 2 months to 60 years old, with Type 1, 2, or 3 SMA



Evrysdi is the first and only oral treatment for SMA that is delivered to your door

- Evrysdi is a liquid taken by mouth or feeding tube once a day

Talk with your healthcare provider to learn more and find out if Evrysdi is right for you

What is Evrysdi?

Evrysdi is a prescription medicine used to treat spinal muscular atrophy (SMA) in adults and children 2 months of age and older.

It is not known if Evrysdi is safe and effective in children under 2 months of age.

Important Safety Information

- Before taking Evrysdi, tell your doctor if you have liver problems, are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed. Evrysdi may harm an unborn or breastfed baby. Evrysdi may affect a man's ability to have children (fertility). Tell your doctor about all the medicines you take.

These are not all the possible side effects of Evrysdi.

Please see additional Important Safety Information and full Prescribing Information.

Genentech
A Member of the Roche Group

© 2020 Genentech USA, Inc. All rights reserved M-US-00006222(v1.0) 08/2020



About
Evrysdi

How Evrysdi
was studied

Results in
Type 1 SMA

Results in
Type 2 & 3
SMA

Safety
studied

How to take
Evrysdi

MySMA
Support

Talking to
your doctor

Summary

Important
Safety
Information

Prescribing
Information

Important Safety Information

What is Evrysdi?

Evrysdi is a prescription medicine used to treat spinal muscular atrophy (SMA) in adults and children 2 months of age and older.

It is not known if Evrysdi is safe and effective in children under 2 months of age.

Important Safety Information

- Before taking Evrysdi, tell your healthcare provider about all of your medical conditions, including if you:
 - have liver problems
 - are pregnant or plan to become pregnant. If you are pregnant, or are planning to become pregnant, ask your healthcare provider for advice before taking this medicine. Evrysdi may harm your unborn baby.
 - are a woman who can become pregnant:
 - Before you start your treatment with Evrysdi, your healthcare provider may test you for pregnancy. Because Evrysdi may harm your unborn baby, your healthcare provider will decide if taking Evrysdi is right for you during this time
 - Talk to your healthcare provider about birth control methods that may be right for you. Use birth control while on treatment and for at least 1 month after stopping Evrysdi
 - are an adult male planning to have children: Evrysdi may affect a man’s ability to have children (fertility). If this is of concern to you, make sure to ask a healthcare provider for advice
 - are breastfeeding or plan to breastfeed. It is not known if Evrysdi passes into breast milk and may harm your baby. If you plan to breastfeed, discuss with your healthcare provider about the best way to feed your baby while on treatment with Evrysdi
- **Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine
- You should receive Evrysdi from the pharmacy as a liquid that can be given by mouth or through a feeding tube. The liquid solution is prepared by your pharmacist. If the medicine in the bottle is a powder, **do not use it**. Contact your pharmacist for a replacement
- Avoid getting Evrysdi on your skin or in your eyes. If Evrysdi gets on your skin, wash the area with soap and water. If Evrysdi gets in your eyes, rinse your eyes with water
- The most common side effects of Evrysdi include:
 - For later-onset SMA:
 - fever
 - diarrhea
 - rash
 - For infantile-onset SMA:
 - fever
 - diarrhea
 - rash
 - runny nose, sneezing, sore throat, and cough (upper respiratory infection)
 - lung infection
 - constipation
 - vomiting

These are not all of the possible side effects of Evrysdi. For more information on the risk and benefits profile of Evrysdi, ask your healthcare provider or pharmacist.

You may report side effects to the FDA at **1-800-FDA-1088** or **www.fda.gov/medwatch**. You may also report side effects to Genentech at **1-888-835-2555**.

Please see full Prescribing Information for additional Important Safety Information.



HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use EVRYSdi safely and effectively. See full prescribing information for EVRYSdi.

EVRYSdi™ (risdiplam) for oral solution
Initial U.S. Approval: 2020

INDICATIONS AND USAGE

EVRYSdi is a survival of motor neuron 2 (SMN2) splicing modifier indicated for the treatment of spinal muscular atrophy (SMA) in patients 2 months of age and older. (1)

DOSAGE AND ADMINISTRATION

EVRYSdi must be constituted by a pharmacist prior to dispensing. Administer orally once daily after a meal using the provided oral syringe. (2.1, 2.4)

Age and Body Weight	Recommended Daily Dosage
2 months to less than 2 years of age	0.2 mg/kg
2 years of age and older weighing less than 20 kg	0.25 mg/kg
2 years of age and older weighing 20 kg or more	5 mg

See Full Prescribing Information for important preparation and administration instructions. (2.1, 2.4)

DOSAGE FORMS AND STRENGTHS

For Oral Solution: 60 mg of risdiplam as a powder for constitution to provide 0.75 mg/mL solution. (3)

CONTRAINDICATIONS

None. (4)

ADVERSE REACTIONS

The most common adverse reactions in later-onset SMA (incidence at least 10% of patients treated with EVRYSdi and more frequent than control) were fever, diarrhea, and rash. (6.1)

The most common adverse reactions in infantile-onset SMA were similar to those observed in later-onset SMA patients. Additionally, adverse reactions with an incidence of at least 10% were upper respiratory tract infection, pneumonia, constipation, and vomiting. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Genentech at 1-888-835-2555 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

Avoid coadministration with drugs that are substrates of multidrug and toxin extrusion (MATE) transporters. (7.1)

USE IN SPECIFIC POPULATIONS

Pregnancy: Based on animal data, may cause fetal harm. (8.1)
Hepatic Impairment: Avoid use in patients with hepatic impairment. (8.6)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 8/2020

FULL PRESCRIBING INFORMATION: CONTENTS*

- 1 INDICATIONS AND USAGE
- 2 DOSAGE AND ADMINISTRATION

2.1 Important Administration Instructions

2.2 Dosing Information

2.3 Missed Dose

2.4 Preparation of Oral Solution by Pharmacists
- 3 DOSAGE FORMS AND STRENGTHS
- 4 CONTRAINDICATIONS
- 6 ADVERSE REACTIONS

6.1 Clinical Trials Experience
- 7 DRUG INTERACTIONS

7.1 Effect of EVRYSdi on Substrates of Multidrug and Toxin Extrusion (MATE) Protein Transporters
- 8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.3 Females and Males of Reproductive Potential

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Hepatic Impairment

- 11 DESCRIPTION
- 12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

13.2 Animal Toxicology and/or Pharmacology
- 14 CLINICAL STUDIES

14.1 Infantile-Onset SMA

14.2 Later-Onset SMA
- 16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

16.2 Storage and Handling
- 17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

Reference ID: 4653194



About
Evrysdi

How Evrysdi
was studied

Results in
Type 1 SMA

Results in
Type 2 & 3
SMA

Safety
studied

How to take
Evrysdi

MySMA
Support

Talking to
your doctor

Summary

Important
Safety
Information

Prescribing
Information

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

EVRYSDI is indicated for the treatment of spinal muscular atrophy (SMA) in patients 2 months of age and older.

2 DOSAGE AND ADMINISTRATION

2.1 Important Administration Instructions

Dose Preparation

It is recommended that a healthcare provider discuss with the patient or caregiver how to prepare the prescribed daily dose prior to administration of the first dose [see Instructions for Use].

Instruct patients or caregivers to prepare the dose using the reusable oral syringe provided.

EVRYSDI must be taken immediately after it is drawn up into the oral syringe. If EVRYSDI is not taken within 5 minutes, EVRYSDI should be discarded from the oral syringe, and a new dose should be prepared.

Dose Administration

EVRYSDI is taken orally once daily after a meal at approximately the same time each day.

In infants who are breastfed, EVRYSDI should be administered after breastfeeding. EVRYSDI cannot be mixed with formula or milk.

Instruct patients to drink water after taking EVRYSDI to ensure the drug has been completely swallowed.

If the patient is unable to swallow and has a nasogastric or gastrostomy tube, EVRYSDI can be administered via the tube. The tube should be flushed with water after delivering EVRYSDI [see Instructions for Use].

2.2 Dosing Information

EVRYSDI is administered orally once daily. The recommended dosage is determined by age and body weight (see Table 1).

Table 1 Adult and Pediatric Dosing Regimen by Age and Body Weight

Age and Body Weight	Recommended Daily Dosage
2 months to less than 2 years of age	0.2 mg/kg
2 years of age and older weighing less than 20 kg	0.25 mg/kg
2 years of age and older weighing 20 kg or more	5 mg

2.3 Missed Dose

If a dose of EVRYSDI is missed, EVRYSDI should be administered as soon as possible if still within 6 hours of the missed dose, and the usual dosing schedule can be resumed on the next day. Otherwise, the missed dose should be skipped, and the next dose should be taken at the regularly scheduled time on the next day.

If a dose is not fully swallowed or vomiting occurs after taking a dose of EVRYSDI, another dose should not be administered to make up for the lost dose. The patient should wait until the next day to take the next dose at the regularly scheduled time.

Reference ID: 4653194



2.4 Preparation of Oral Solution by Pharmacists

EVRYSDI powder must be constituted to the oral solution by a pharmacist prior to dispensing to the patient.

Preparation of the EVRYSDI Oral Solution 0.75 mg/mL

The EVRYSDI “Instructions for Constitution” booklet contains more detailed instructions on the preparation of the oral solution [see *Instructions for Constitution*].

Caution should be exercised in the handling of EVRYSDI powder for oral solution. Avoid inhalation and direct contact with skin or mucous membranes with the dry powder and the constituted solution. If such contact occurs, wash thoroughly with soap and water; rinse eyes with water. Wear disposable gloves during the preparation and cleanup procedure.

1. Gently tap the bottom of the closed glass bottle to loosen the powder.
2. Remove the cap. Do not throw away the cap.
3. Carefully pour 79 mL of Purified Water into the EVRYSDI bottle to yield the 0.75 mg/mL oral solution. Do not mix EVRYSDI with formula or milk.
4. Insert the Press-In bottle adapter into the bottle opening by pushing it down against the bottle lip. Ensure it is completely pressed against the bottle lip.
5. Re-cap the bottle tightly and shake well for 15 seconds. Wait for 10 minutes. You should have obtained a clear solution. If not, shake well again for another 15 seconds.
6. Write the date of expiration of the constituted oral solution (calculated as 64 days after constitution) and the lot number on the bottle label. Peel off the part of the bottle label that has the expiration date of the powder.
7. Put the bottle back in its original carton.
8. Select the appropriate oral syringes (6 mL or 12 mL) based on the patient’s dosage and remove the other oral syringes from the carton.
9. Dispense with the “Instructions for Use” and FDA-approved patient labeling. Alert patients to read the important handling information described in the Instructions for Use.

Keep the constituted oral solution of EVRYSDI in the original amber bottle to protect from light. Store in a refrigerator at 2°C to 8°C (36°F to 46°F). Do not freeze. Discard any unused portion 64 days after constitution. Keep the bottle in an upright position with the cap tightly closed.

3 DOSAGE FORMS AND STRENGTHS

EVRYSDI for oral solution: 60 mg as a light yellow, yellow, greyish yellow, greenish yellow, or light green powder for constitution. Following constitution, the volume of the greenish yellow to yellow solution is 80 mL, providing 60 mg/80 mL (0.75 mg/mL) risdiplam.

4 CONTRAINDICATIONS

None.

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in clinical trials of another drug and may not reflect the rates observed in practice.

Reference ID: 4653194



In clinical trials including patients with infantile-onset SMA and later-onset SMA, a total of 337 patients (52% female, 72% Caucasian) were exposed to EVRYSDI for up to a maximum of 32 months, with 209 patients receiving treatment for more than 12 months. Forty-seven (14%) patients were 18 years and older, 74 (22%) were 12 years to less than 18 years, 154 (46%) were 2 years to less than 12 years, and 62 (18%) 2 months to less than 2 years.

Clinical Trial in Later-Onset SMA

The safety of EVRYSDI for later-onset SMA is based on data from a randomized, double-blinded, placebo-controlled study (Study 2 Part 2) in patients with SMA Type 2 or 3 (n = 180) [see Clinical Studies (14.2)]. The patient population in Study 2 Part 2 ranged in age from 2 to 25 years at the time of treatment start.

The most common adverse reactions (reported in at least 10% of patients treated with EVRYSDI and at an incidence greater than on placebo) in Study 2 Part 2 were fever, diarrhea, and rash. Table 2 lists the adverse reactions that occurred in at least 5% of patients treated with EVRYSDI and at an incidence ≥ 5% greater than on placebo in Study 2 Part 2.

Table 2 Adverse Reactions Reported in ≥ 5% of Patients Treated with EVRYSDI and with an Incidence ≥ 5% Greater Than on Placebo in Study 2 Part 2

Adverse Reaction	EVRYSDI (N = 120) %	Placebo (N = 60) %
Fever ¹	22	17
Diarrhea	17	8
Rash ²	17	2
Mouth and aphthous ulcers	7	0
Arthralgia	5	0
Urinary tract infection ³	5	0

¹ Includes pyrexia and hyperpyrexia.
² Includes rash, erythema, rash maculo-papular, rash erythematous, rash papular, dermatitis allergic, and folliculitis.
³ Includes urinary tract infection and cystitis.

Clinical Trial in Infantile-Onset SMA

The safety of EVRYSDI therapy for infantile-onset SMA is based on data from an open-label study in 62 patients (Study 1) [see Clinical Studies (14.1)]. In Study 1 Part 1 (n = 21) and Part 2 (n = 41), 62 patients received EVRYSDI for up to 30 months (31 patients for more than 12 months). The patient population ranged in age from 2 to 7 months at the time of treatment start (weight range 4.1 to 10.6 kg).

The most frequent adverse reactions reported in infantile-onset SMA patients treated with EVRYSDI in Study 1 were similar to those observed in later-onset SMA patients in Study 2. Additionally, the following adverse reactions were reported in ≥ 10% of patients: upper respiratory tract infection (including nasopharyngitis, rhinitis, respiratory tract infection), pneumonia, constipation, and vomiting.

Reference ID: 4653194



7 DRUG INTERACTIONS

7.1 Effect of EVRYSdi on Substrates of Multidrug and Toxin Extrusion (MATE) Protein Transporters

Based on in vitro data, EVRYSdi may increase plasma concentrations of drugs eliminated via MATE1 or MATE2-K [see Clinical Pharmacology (12.3)], such as metformin. Avoid coadministration of EVRYSdi with MATE substrates. If coadministration cannot be avoided, monitor for drug-related toxicities and consider dosage reduction of the coadministered drug (based on the labeling of that drug) if needed.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no adequate data on the developmental risk associated with the use of EVRYSdi in pregnant women. In animal studies, administration of risdiplam during pregnancy or throughout pregnancy and lactation resulted in adverse effects on development (embryofetal mortality, malformations, decreased fetal body weights, and reproductive impairment in offspring) at or above clinically relevant drug exposures [see Data].

The estimated background risk of major birth defects and miscarriage for the indicated populations is unknown. In the U.S. general population, the estimated background risk of major birth defect and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively. Based on animal data, advise pregnant women of the potential risk to the fetus.

Data

Animal Data

Oral administration of risdiplam (0, 1, 3, or 7.5 mg/kg) to pregnant rats throughout organogenesis resulted in decreased fetal body weights and increased incidences of fetal structural variations at the highest dose tested, which was not associated with maternal toxicity. The no-effect level for adverse effects on embryofetal development (3 mg/kg/day) was associated with maternal plasma exposure (AUC) approximately 2 times that in humans at the maximum recommended human dose (MRHD) of 5 mg.

Oral administration of risdiplam (0, 1, 4, or 12 mg/kg) to pregnant rabbits throughout organogenesis resulted in embryofetal mortality, fetal malformations (hydrocephaly), and structural variations at the highest dose tested, which was associated with maternal toxicity. The no-effect dose for adverse effects on embryofetal development (4 mg/kg/day) was associated with maternal plasma exposure (AUC) approximately 4 times that in humans at the MRHD.

When risdiplam (0, 0.75, 1.5, or 3 mg/kg/day) was orally administered to rats throughout pregnancy and lactation, gestation was prolonged in the dams, and delayed sexual maturation (vaginal opening) and impaired reproductive function (decreased numbers of corpora lutea, implantation sites, and live embryos) were observed in female offspring at the highest dose. The no-effect dose for adverse effects on pre- and postnatal development in rats (1.5 mg/kg/day) was associated with maternal plasma exposure (AUC) similar to that in humans at the MRHD.

8.2 Lactation

Risk Summary

There are no data on the presence of risdiplam in human milk, the effects on the breastfed infant, or the effects on milk production. Risdiplam was excreted in the milk of lactating rats orally administered risdiplam.

Reference ID: 4653194



The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for EVRYSdi and any potential adverse effects on the breastfed infant from EVRYSdi or from the underlying maternal condition.

8.3 Females and Males of Reproductive Potential

Studies of risdiplam in juvenile and adult rats and in monkeys demonstrated adverse effects on the reproductive organs, including germ cells, in males at clinically-relevant plasma exposures [see Use in Specific Populations (8.4) and Nonclinical Toxicology (13.1)].

Pregnancy Testing

Pregnancy testing is recommended for females of reproductive potential prior to initiating EVRYSdi [see Use in Specific Populations (8.1)].

Contraception

EVRYSdi may cause embryofetal harm when administered to a pregnant woman [see Use in Specific Populations (8.1)].

Female Patients

Advise female patients of reproductive potential to use effective contraception during treatment with EVRYSdi and for at least 1 month after her last dose.

Infertility

Male Patients

Male fertility may be compromised by treatment with EVRYSdi [see Nonclinical Toxicology (13.1)].

Counsel male patients of reproductive potential receiving EVRYSdi about the potential effects on fertility. Male patients may consider sperm preservation prior to treatment.

8.4 Pediatric Use

The safety and effectiveness of EVRYSdi in pediatric patients 2 months of age and older have been established [see Clinical Studies (14)]. Safety and effectiveness in pediatric patients below the age of 2 months have not been established [see Clinical Pharmacology (12.3)].

Juvenile Animal Toxicity Data

Oral administration of risdiplam (0, 0.75, 1.5, 2.5 mg/day) to young rats from postnatal day (PND) 4 through PND 31 resulted in decreased growth (body weight, tibia length) and delayed sexual maturation in males at the mid and high dose. The skeletal and body weight deficits persisted after cessation of dosing. Ophthalmic changes consisting of vacuoles in the anterior vitreous were seen at the high dose. Decreases in absolute B lymphocyte counts were observed at all doses after cessation of dosing. Decreases in testis and epididymis weights, which correlated with degeneration of the seminiferous epithelium in the testis, occurred at the mid and high doses; the histopathology findings were reversible, but organ weight persisted after cessation of dosing. Impaired female reproductive performance (decreased mating index, fertility index, and conception rate) was observed at the high dose. A no-effect dose for adverse developmental effects on preweaning rats was not identified. The lowest dose tested (0.75 mg/kg/day) was associated with plasma exposures (AUC) lower than that in humans at the maximum recommended human dose (MRHD) of 5 mg/day.

Oral administration of risdiplam (0, 1, 3, or 7.5 mg/day) to young rats from PND 22 through PND 112 produced a marked increase in micronuclei in the bone marrow, male reproductive organ histopathology (degeneration/necrosis of the seminiferous tubule epithelium,

Reference ID: 4653194



oligo/aspermia in the epididymis, spermatid granulomas), and adverse effects on sperm parameters (decreased sperm concentration and motility, increased sperm morphology abnormalities) at the highest dose tested. Increases in T lymphocytes (total, helper, and cytotoxic) were observed at the mid and high doses. The reproductive and immune effects persisted after cessation of dosing. The no-effect dose (1 mg/kg/day) for adverse effects on postweaning juvenile rats was associated with plasma exposures (AUC) lower than that in humans at the MRHD.

8.5 Geriatric Use

Clinical studies of EVRYSDI did not include patients aged 65 years and over to determine whether they respond differently from younger patients.

8.6 Hepatic Impairment


The safety and efficacy of EVRYSDI in patients with hepatic impairment have not been studied. Because risdiplam is predominantly metabolized in the liver, hepatic impairment may potentially increase the exposures to risdiplam [see *Clinical Pharmacology* (12.3)]. Avoid use of EVRYSDI in patients with impaired hepatic function.

11 DESCRIPTION

EVRYSDI for oral solution contains risdiplam, which is a survival of motor neuron 2 (SMN2)-directed RNA splicing modifier.

The chemical name of risdiplam is 7-(4,7-diazaspiro[2.5]octan-7-yl)-2-(2,8-dimethylimidazo[1,2-b]pyridazin-6-yl)pyrido-4H-[1,2-a]pyrimidin-4-one. Risdiplam has a molecular weight of 401.46 g/mol.

The molecular formula of risdiplam is C₂₂H₂₃N₇O and the chemical structure is shown below.



EVRYSDI is supplied as a powder in an amber glass bottle. Each bottle contains 60 mg of risdiplam. The inactive ingredients of EVRYSDI are: ascorbic acid, disodium edetate dihydrate, isomalt, mannitol, polyethylene glycol 6000, sodium benzoate, strawberry flavor, sucralose, and tartaric acid.

The powder is constituted with purified water to yield 60 mg/80 mL (0.75 mg/mL) of risdiplam after constitution [see *Dosage and Administration* (2.4)].

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Risdiplam is a survival of motor neuron 2 (SMN2) splicing modifier designed to treat patients with spinal muscular atrophy (SMA) caused by mutations in chromosome 5q that lead to SMN protein deficiency. Using in vitro assays and studies in transgenic animal models of SMA, risdiplam was shown to increase exon 7 inclusion in SMN2 messenger ribonucleic acid (mRNA) transcripts and production of full-length SMN protein in the brain.

In vitro and in vivo data indicate that risdiplam may cause alternative splicing of additional genes, including FOXM1 and MADD. FOXM1 and MADD are thought to be involved in cell

Reference ID: 4653194



About
Evrysdi

How Evrysdi
was studied

Results in
Type 1 SMA

Results in
Type 2 & 3
SMA

Safety
studied

How to take
Evrysdi

MySMA
Support

Talking to
your doctor

Summary

Important
Safety
Information

Prescribing
Information

cycle regulation and apoptosis, respectively, and have been identified as possible contributors to adverse effects seen in animals.

12.2 Pharmacodynamics

In clinical trials, EVRYSDI led to an increase in SMN protein with a greater than 2-fold median change from baseline within 4 weeks of treatment initiation. The increase was sustained throughout the treatment period (of at least 12 months) across all SMA types.

12.3 Pharmacokinetics

Pharmacokinetics of EVRYSDI have been characterized in healthy adult subjects and in patients with SMA.

After administration of EVRYSDI as an oral solution, pharmacokinetics of risdiplam were approximately linear between 0.6 and 18 mg in a single-ascending-dose study in healthy adult subjects, and between 0.02 and 0.25 mg/kg once daily in a multiple-ascending-dose study in patients with SMA. Following once-daily oral administration of risdiplam in healthy subjects, approximately 3-fold accumulation of peak plasma concentrations (C_{max}) and area under the plasma concentration-time curve (AUC_{0-24h}) was observed. Risdiplam exposures reach steady state 7 to 14 days after once-daily administration.

Absorption

Following oral administration, the time to reach maximum plasma concentration (T_{max}) is between 1 and 4 hours.

Effect of Food

In the clinical efficacy studies (Study 1 and Study 2), risdiplam was administered with a morning meal or after breastfeeding.

Distribution

The apparent volume of distribution at steady state is 6.3 L/kg.

Risdiplam is predominantly bound to serum albumin, without any binding to alpha-1 acid glycoprotein, with a free fraction of 11%.

Elimination

The apparent clearance (CL/F) of risdiplam is 2.1 L/h for a 14.9 kg patient.

The terminal elimination half-life of risdiplam was approximately 50 hours in healthy adults.

Metabolism

Risdiplam is primarily metabolized by flavin monooxygenase 1 and 3 (FMO1 and FMO3) and also by CYPs 1A1, 2J2, 3A4, and 3A7.

Parent drug was the major component found in plasma, accounting for 83% of drug-related material in circulation. The pharmacologically-inactive metabolite M1 was identified as the major circulating metabolite.

Excretion

Following a dose of 18 mg, approximately 53% of the dose (14% unchanged risdiplam) was excreted in the feces and 28% in urine (8% unchanged risdiplam).

Specific Populations

There were no clinically significant differences in the pharmacokinetics of EVRYSDI based on race or gender. Renal impairment is not expected to alter the exposures to risdiplam.

Reference ID: 4653194



The impact of geriatric age and hepatic impairment on the pharmacokinetics of EVRYSDI has not been studied.

Pediatric Patients

Body weight and age were found to have significant effect on the pharmacokinetics of risdiplam. The estimated exposure (mean AUC_{0-24h}) for infantile-onset SMA patients (age 2 to 7 months at enrollment) at the recommended dose of 0.2 mg/kg once daily was 1930 ng.h/mL. The estimated exposure for later-onset SMA patients (2 to 25 years old at enrollment) at the recommended dose was 2050 ng.h/mL (0.25 mg/kg once daily for patients with a body weight < 20 kg and 5 mg once daily for patients with a body weight ≥ 20 kg). The observed maximum concentration (mean C_{max}) was 184 ng/mL for infantile-onset SMA patients and 148 ng/mL for later-onset SMA patients.

Based on literature reports, pediatric patients less than 2 months of age are expected to have reduced activity of FMO3, which may result in increased exposure to risdiplam [see *Elimination*]. No data on risdiplam pharmacokinetics are available in patients less than 2 months of age [see *Use in Specific Populations (8.4)*].

Drug Interaction Studies

Effect of Other Drugs on EVRYSDI

Coadministration of 200 mg itraconazole (a strong CYP3A inhibitor) twice daily with a single 6 mg oral dose of risdiplam did not have a clinically relevant effect on the pharmacokinetics of risdiplam (11% increase in AUC and 9% decrease in C_{max}).

Risdiplam is a weak substrate of human MDR-1 and breast cancer resistant protein (BCRP) transporters in vitro. Human MDR-1 or BCRP inhibitors are not expected to result in a clinically significant increase of risdiplam concentrations.

Effect of EVRYSDI on Other Drugs

Risdiplam and its major circulating metabolite M1 did not induce CYP1A2, 2B6, 2C8, 2C9, 2C19, or 3A4 in vitro. Risdiplam and M1 did not inhibit (reversible or time-dependent inhibition) any of the CYP enzymes tested (CYP1A2, 2B6, 2C8, 2C9, 2C19, 2D6), with the exception of CYP3A in vitro.

EVRYSDI is a weak inhibitor of CYP3A. In healthy adult subjects, administration of EVRYSDI once daily for 2 weeks slightly increased the exposure of midazolam, a sensitive CYP3A substrate (AUC 11%; C_{max} 16%); this increase is not considered clinically relevant. Based on physiologically-based pharmacokinetic (PBPK) modeling, a similar increase is expected in children and infants as young as 2 months of age.

In vitro studies have shown that risdiplam and its major metabolite are not significant inhibitors of human MDR1, organic anion-transporting polypeptide (OATP) 1B1, OATP1B3, organic anion transporter 1 and 3 (OAT 1 and 3) transporters, and human organic cation transporter 2 (OCT2), at clinically relevant concentrations. Risdiplam and its metabolite are, however, in vitro inhibitors of the multidrug and toxin extrusion (MATE) 1 and MATE2-K transporters [see *Drug Interactions (7.1)*].

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Reference ID: 4653194



About
Evrysdi

How Evrysdi
was studied

Results in
Type 1 SMA

Results in
Type 2 & 3
SMA

Safety
studied

How to take
Evrysdi

MySMA
Support

Talking to
your doctor

Summary

Important
Safety
Information

Prescribing
Information

The carcinogenic potential of risdiplam has not been fully evaluated. Risdiplam was not carcinogenic in Tg.rasH2 mice when administered at oral doses of up to 9 mg/kg/day for 26 weeks.

Mutagenesis

Risdiplam was negative in an in vitro Ames assay. In an in vivo combined bone marrow micronucleus and comet assay in rat, risdiplam was clastogenic, as evidenced by an increase in micronuclei in bone marrow, but was negative in the comet assay. A pronounced increase in bone marrow micronuclei was also observed in toxicity studies in adult and juvenile rats [see *Use in Specific Populations (8.4)*].

Impairment of Fertility

Oral administration of risdiplam to rats for 4 (0, 1, 3, or 9 mg/kg/day) or 26 (0, 1, 3, or 7.5 mg/kg/day) weeks resulted in histopathological effects in the testis (degenerated spermatocytes, degeneration/atrophy of the seminiferous tubules) and epididymis (degeneration/necrosis of ductular epithelium) at the mid and/or high doses. At the high dose in the 26-week study, the testicular lesions persisted to the end of the recovery period, which corresponds, in rat, to approximately one spermatogenic cycle. The no-effect dose for adverse reproductive system effects in adult male rats (1 mg/kg/day) was associated with plasma drug exposures (AUC) similar to that in humans at the maximum recommended human dose (MRHD) of 5 mg/day.

Adverse effects of risdiplam on the testis could not be fully evaluated in the monkey because the majority of monkeys tested were sexually immature. However, oral administration of risdiplam (0, 2, 4, or 6 mg/kg/day) for 2 weeks resulted in histopathological changes in the testis (increases in multinucleate cells, germ cell degeneration) at the highest dose. At the no-effect dose for testicular toxicity in monkeys, plasma exposures were approximately 3 times that in humans at the MRHD.

Oral administration of risdiplam to postweaning juvenile rats resulted in male reproductive toxicity (degeneration/necrosis of the testis seminiferous epithelium with associated oligo/azpermia in the epididymis and abnormal sperm parameters). The no-effect dose for adverse reproductive effects in postweaning male juvenile rats was associated with plasma exposures approximately 4 times that in humans at the MRHD [see *Use in Specific Populations (8.4)*].

13.2 Animal Toxicology and/or Pharmacology

Retinal toxicity

Risdiplam-induced functional and structural retinal abnormalities were seen in animal studies. In a 39-week toxicity study in monkeys, oral administration of risdiplam (0, 1.5, 3, or 7.5/5 mg/kg/day; high dose lowered after 4 weeks) produced functional abnormalities on the electroretinogram (ERG) in all mid- and high-dose animals at the earliest examination time (Week 20). These findings were associated with retinal degeneration, detected by optical coherence tomography (OCT), on Week 22, the first examination time. The retinal degeneration, with peripheral photoreceptor loss, was irreversible. A no-effect dose for the retinal findings (1.5 mg/kg/day) was associated with plasma exposures (AUC) similar to that in humans at the maximum recommended human dose (MRHD) of 5 mg.

Effect on Epithelial Tissues

Oral administration of risdiplam to rats and monkeys resulted in histopathological changes in epithelium of the gastrointestinal (GI) tract (apoptosis/single cell necrosis), lamina propria (vacuolation), the exocrine pancreas (single cell necrosis), the skin, tongue, and larynx

Reference ID: 4653194



(parakeratosis/hyperplasia/degeneration) with associated inflammation. The skin and GI epithelial effects were reversible. The no-effect doses for effects on epithelial tissues in rats and monkeys were associated with plasma exposures (AUC) similar to that in humans at the MRHD.

14 CLINICAL STUDIES

The efficacy of EVRYSDI for the treatment of patients with infantile-onset and later-onset SMA was evaluated in two clinical studies, Study 1 (NCT02913482) and Study 2 (NCT02908685).

The overall findings of these studies support the effectiveness of EVRYSDI in SMA patients 2 months of age and older and appear to support the early initiation of treatment with EVRYSDI.

14.1 Infantile-Onset SMA

Study 1 was an open-label, 2-part study to investigate the efficacy, safety, pharmacokinetics, and pharmacodynamics of EVRYSDI in patients with Type 1 SMA (symptom onset between 28 days and 3 months of age). Part 1 of Study 1 (n = 21) provides efficacy and safety data in patients with Type 1 SMA. Additional safety information is provided by Part 2 of Study 1 (n = 41) in patients with Type 1 SMA [see Adverse Reactions (6.1)].

In Part 1 of Study 1, patients (n = 21) were enrolled in one of two dosage cohorts. Patients in the higher-dosage cohort (n = 17) had their dosage adjusted to the recommended dosage of 0.2 mg/kg/day before 12 months of treatment, while patients in the low-dosage cohort (n = 4) did not.

Effectiveness was established based on the ability to sit without support for at least 5 seconds (as measured by Item 22 of the Bayley Scales of Infant and Toddler Development – Third Edition (BSID-III) gross motor scale) and on the basis of survival without permanent ventilation. Permanent ventilation was defined as requiring a tracheostomy or more than 21 consecutive days of either non-invasive ventilation (≥ 16 hours per day) or intubation, in the absence of an acute reversible event.

The median age of onset of clinical signs and symptoms of Type 1 SMA in patients enrolled in Part 1 of Study 1 was 2.0 months (range: 0.9 to 3.0); 71% of patients were female, 81% were Caucasian, and 19% were Asian. The median age at enrollment was 6.7 months (range: 3.3 to 6.9), and the median time between onset of symptoms and first dose was 4.0 months (range: 2.0 to 5.8). All patients had genetic confirmation of homozygous deletion or compound heterozygosity predictive of loss of function of the SMN1 gene, and two SMN2 gene copies.

In Study 1 Part 1, the median duration of EVRYSDI treatment was 14.8 months (range: 0.6 to 26.0), and 19 patients were treated for a minimum duration of 12 months.

Of the patients who were treated with the recommended dosage of EVRYSDI 0.2 mg/kg/day, 41% (7/17) were able to sit independently for ≥ 5 seconds (BSID-III, Item 22) after 12 months of treatment. These results indicate a clinically meaningful deviation from the natural history of untreated infantile-onset SMA. As described in the natural history of untreated infantile-onset SMA, patients would not be expected to attain the ability to sit independently, and no more than 25% of these patients would be expected to survive without permanent ventilation beyond 14 months of age. After 12 months of treatment with EVRYSDI, 90% (19/21) of patients were alive without permanent ventilation (and reached 15 months of age or older). After a minimum of 23 months of treatment with EVRYSDI, 81% (17/21) of all patients were alive without permanent ventilation (and reached an age of 28 months or older; median 32 months; range 28 to 45 months).

14.2 Later-Onset SMA

Study 2 was a 2-part, multicenter trial to investigate the efficacy, safety, pharmacokinetics, and pharmacodynamics of EVRYSDI in patients diagnosed with SMA Type 2 or Type 3. Part 1 of

Reference ID: 4653194



Study 2 was dose-finding and exploratory in 51 patients (14% ambulatory). Part 2 was randomized, double-blind, placebo-controlled, and is described below.

The primary endpoint in Study 2 Part 2 was the change from baseline to Month 12 in the Motor Function Measure 32 (MFM32) score. A key secondary endpoint was the proportion of patients with a 3-point or greater change from baseline to Month 12 in the MFM32 total score. The MFM32 measures motor function abilities that relate to daily functions. The total MFM32 score is expressed as a percentage (range: 0 to 100) of the maximum possible score, with higher scores indicating greater motor function. Another key secondary endpoint was the Revised Upper Limb Module (RULM). The RULM is a tool used to assess motor performance of the upper limb in SMA patients. It tests proximal and distal motor functions of the arm. The total score ranges from 0 (all the items cannot be performed) to 37 (all the activities are achieved fully without any compensatory maneuvers).

Study 2 Part 2 enrolled 180 non-ambulatory patients with Type 2 (71%) or Type 3 (29%) SMA. Patients were randomized 2:1 to receive EVRYSdi at the recommended dosage [see *Dosage and Administration* (2.2)] or placebo. Randomization was stratified by age group (2 to 5, 6 to 11, 12 to 17, or 18 to 25 years of age).

The median age of patients at the start of treatment was 9.0 years (range 2 to 25), and the median time between onset of initial SMA symptoms and first treatment was 102.6 months (range 1 to 275). Of the 180 patients included in the trial, 51% were female, 67% were Caucasian, and 19% were Asian. At baseline, 67% of patients had scoliosis (32% of them with severe scoliosis). Patients had a mean baseline MFM32 score of 46.1, and RULM score of 20.1. Overall baseline demographic characteristics were reasonably balanced between the treatment groups (EVRYSdi and placebo), with the exception of scoliosis (63% in the EVRYSdi arm vs. 73% in the placebo group).

The primary analysis on the change from baseline in MFM32 total score at Month 12 showed a clinically meaningful and statistically significant difference between patients treated with EVRYSdi and placebo. The results of the primary analysis and key secondary endpoints are shown in Table 3 and Figure 1.

Reference ID: 4653194



Table 3 Summary of Efficacy in Patients with Later-Onset SMA at Month 12 of Treatment (Study 2 Part 2)

Endpoint	EVRYSDI (N = 120)	Placebo (N = 60)
Primary Endpoint:		
Change from baseline in total MFM32 score at Month 12, LS means (95% CI) ^{1,2,3}	1.36 (0.61, 2.11)	-0.19 (-1.22, 0.84)
Difference from Placebo, Estimate (95% CI) ¹ p-value	1.55 (0.30, 2.81) 0.0156	
Secondary Endpoints:		
Proportion of patients with a change from baseline MFM32 total score of 3 or more at Month 12 (95% CI) ^{2,3}	38.3% (28.9, 47.6)	23.7% (12.0, 35.4)
Odds ratio for overall response (95% CI) adjusted ⁴ (unadjusted) p-value ⁵	2.35 (1.01, 5.44) 0.0469 (0.0469)	
Change from baseline in total score of RULM at Month 12, LS means (95% CI) ^{1, 6}	1.61 (1.00, 2.22)	0.02 (-0.83, 0.87)
Difference from Placebo, Estimate (95% CI) adjusted ⁴ (unadjusted) p-value ¹	1.59 (0.55, 2.62) 0.0469 (0.0028)	

^{1.} The Mixed Model Repeated Measure (MMRM) analysis included the change from baseline total score as the dependent variable and as independent variables the baseline total score, treatment group, time, treatment-by-time interaction, and the randomization stratification variable of age group (2 to 5, 6 to 11, 12 to 17, 18 to 25).

^{2.} The MFM total score was calculated according to the user manual, expressed as a percentage of the maximum score possible for the scale (i.e., sum of the 32 item scores divided by 96 and multiplied by 100).

^{3.} Based on the missing data rule for MFM32, 6 patients were excluded from the analysis (EVRYSDI n = 115; placebo control n = 59).

^{4.} The adjusted p-value was obtained for the endpoints included in the hierarchical testing and was derived based on all the p-values from endpoints in order of the hierarchy up to the current endpoint.


^{5.} The logistic regression analysis included the baseline total score, treatment and age group as independent variables.

^{6.} Based on the missing data rule for RULM, 3 patients were excluded from the analysis (EVRYSDI n = 119; placebo control n = 58).

Reference ID: 4653194



Figure 1 Mean Change from Baseline in Total MFM32 Score Over 12 Months (Study 2 Part 2)^{1,2}



¹ Error bars denote the 95% confidence interval.
² The MFM total score was calculated according to the user manual, expressed as a percentage of the maximum score possible for the scale (i.e., sum of the 32 item scores divided by 96 and multiplied by 100).

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

Each amber glass bottle of EVRYSDI is packaged with a bottle adapter, two 6 mL reusable oral syringes, and two 12 mL reusable oral syringes. EVRYSDI for oral solution is a light yellow, yellow, greyish yellow, greenish yellow, or light green powder. Each bottle contains 60 mg of risdiplam (NDC 50242-175-07).

16.2 Storage and Handling

Store the dry powder at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C to 30°C (59°F to 86°F) [see USP controlled room temperature]. Keep in the original carton.
Keep the constituted oral solution of EVRYSDI in the original amber bottle to protect from light. Store in a refrigerator at 2°C to 8°C (36°F to 46°F) [see Dosage and Administration (2.4)].

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information and Instructions for Use).

Pregnancy and Fetal Risk

Inform pregnant women and women of reproductive potential that, based on animal studies, EVRYSDI may cause fetal harm [see Use in Specific Populations (8.1)].
Discuss with women of childbearing age whether they are pregnant, might be pregnant, or are trying to become pregnant.
Advise women of childbearing potential to use effective contraception during treatment with EVRYSDI and for at least 1 month after stopping EVRYSDI.

Reference ID: 4653194



Advise a female patient to immediately inform the prescriber if she is pregnant or planning to become pregnant [see Use in Specific Populations (8.3)].

Potential Effects on Male Fertility

Advise male patients that their fertility may be compromised while on treatment with EVRYSDI [see Use in Specific Populations (8.3)].

Instructions for Preparation of Oral Solution

Advise patients to ensure that EVRYSDI is in liquid form when received from the pharmacy.

Instruct patients/caregivers to take EVRYSDI after a meal or after breastfeeding at approximately the same time each day. However, instruct caregivers to not mix EVRYSDI with formula or milk.

Instruct patients/caregivers to take EVRYSDI immediately after it is drawn up into the reusable oral syringe [see Dosage and Administration (2.1)].

EVRYSDI™ [risdiplam]

Distributed by:

Genentech, Inc.

A Member of the Roche Group

1 DNA Way

South San Francisco, CA 94080-4990

EVRYSDI is a trademark of Genentech, Inc.

©2020 Genentech, Inc. All rights reserved.

Reference ID: 4653194





Patient Information EVRYSDI™ [ev-RIZ-dee] (risdiplam) for oral solution
<p>What is EVRYSDI?</p> <ul style="list-style-type: none">EVRYSDI is a prescription medicine used to treat spinal muscular atrophy (SMA) in adults and children 2 months of age and older.It is not known if EVRYSDI is safe and effective in children under 2 months of age.
<p>Before taking EVRYSDI, tell your healthcare provider about all of your medical conditions, including if you:</p> <ul style="list-style-type: none">have liver problems.are pregnant or plan to become pregnant. If you are pregnant, or are planning to become pregnant, ask your healthcare provider for advice before taking this medicine. EVRYSDI may harm your unborn baby.are a woman who can become pregnant:<ul style="list-style-type: none">Before you start your treatment with EVRYSDI, your healthcare provider may test you for pregnancy. Because EVRYSDI may harm your unborn baby, you and your healthcare provider will decide if taking EVRYSDI is right for you during this time.Talk to your healthcare provider about birth control methods that may be right for you. Use birth control while on treatment and for at least 1 month after stopping EVRYSDI.are an adult male planning to have children: EVRYSDI may affect a man's ability to have children (fertility). If this is of concern to you, make sure to ask a healthcare provider for advice.are breastfeeding or plan to breastfeed. It is not known if EVRYSDI passes into breast milk and may harm your baby. If you plan to breastfeed, discuss with your healthcare provider about the best way to feed your baby while on treatment with EVRYSDI.
<p>Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.</p>
<p>How should I take EVRYSDI?</p> <p>See the detailed Instructions for Use that comes with EVRYSDI for information on how to take or give EVRYSDI oral solution.</p> <ul style="list-style-type: none">You should receive EVRYSDI from the pharmacy as a liquid that can be given by mouth or through a feeding tube. The liquid solution is prepared by your pharmacist. If the medicine in the bottle is a powder, do not use it. Contact your pharmacist for a replacement.Avoid getting EVRYSDI on your skin or in your eyes. If EVRYSDI gets on your skin, wash the area with soap and water. If EVRYSDI gets in your eyes, rinse your eyes with water. <p><i>Taking EVRYSDI</i></p> <ul style="list-style-type: none">Your healthcare provider will tell you how long you or your child needs to take EVRYSDI. Do not stop treatment with EVRYSDI unless your healthcare provider tells you to.For infants and children, your healthcare provider will determine the daily dose of EVRYSDI needed based on your child's age and weight. For adults, take 5 mg of EVRYSDI daily.<ul style="list-style-type: none">Take EVRYSDI exactly as your healthcare provider tells you to take it. Do not change the dose without talking to your healthcare provider.Take EVRYSDI 1 time daily after a meal (or after breastfeeding for a child) at approximately the same time each day. Drink water afterwards to make sure EVRYSDI has been completely swallowed.Do not mix EVRYSDI with formula or milk.If you are unable to swallow and have a nasogastric or gastrostomy tube, EVRYSDI can be given through the tube.If you miss a dose of EVRYSDI:<ul style="list-style-type: none">If you remember the missed dose within 6 hours of when you normally take EVRYSDI, then take or give the dose. Continue taking EVRYSDI at your usual time the next day.If you remember the missed dose more than 6 hours after you normally take EVRYSDI, skip the missed dose. Take your next dose at your usual time the next day.If you do not fully swallow the dose, or you vomit after taking a dose, do not take another dose of EVRYSDI to make up for that dose. Wait until the next day to take the next dose at your usual time.

Reference ID: 4653194





Reusable Oral Syringes

- Your pharmacist will provide you with the reusable oral syringes that are needed for taking your medicine and explain how to use them. Wash the syringes per instructions after use. Do not throw them away.
- Use the reusable oral syringes provided by your pharmacist (you should receive 2 identical oral syringes) to measure your or your child’s dose of EVRYSDI, as they are designed to protect the medicine from light. Contact your healthcare provider or pharmacist if your oral syringes are lost or damaged.
- Once transferred from the bottle to the oral syringe, take EVRYSDI right away. Do not store the EVRYSDI solution in the syringe. If EVRYSDI is not taken within 5 minutes of when it is drawn up, EVRYSDI should be thrown away from the reusable oral syringe, and a new dose should be prepared.

What are the possible side effects of EVRYSDI?

The most common side effects of EVRYSDI include:

- **For later-onset SMA:**
 - fever
 - diarrhea
 - rash
- **For infantile-onset SMA:**
 - fever
 - runny nose, sneezing, sore throat, and cough (upper respiratory infection)
 - constipation
 - diarrhea
 - lung infection
 - vomiting
 - rash

These are not all of the possible side effects of EVRYSDI. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store EVRYSDI?

- Store EVRYSDI in the refrigerator between 36°F to 46°F (2°C to 8°C). Do not freeze.
- Keep EVRYSDI in an upright position in the original amber bottle to protect from light.
- Throw away (discard) any unused portion of EVRYSDI 64 days after it is mixed by the pharmacist (constitution). Please see the Discard After date written on the bottle label. (See the **Instructions for Use** that comes with EVRYSDI).

Keep EVRYSDI and all medicines out of the reach of children.

General information about the safe and effective use of EVRYSDI.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use EVRYSDI for a condition for which it was not prescribed. Do not give EVRYSDI to other people, even if they have the same symptoms you have. It may harm them. You can ask your pharmacist or healthcare provider for information about EVRYSDI that is written for health professionals.

What are the ingredients in EVRYSDI?

Active ingredient: risdiplam

Inactive ingredients: ascorbic acid, disodium edetate dihydrate, isomalt, mannitol, polyethylene glycol 6000, sodium benzoate, strawberry flavor, sucralose, and tartaric acid.

Distributed by: **Genentech, Inc.**, A Member of the Roche Group, 1 DNA Way, South San Francisco, CA 94080-4990
EVRYSDI is a trademark of Genentech, Inc.
©2020 Genentech, Inc. All rights reserved.
For more information, go to www.EVRYSDI.com or call 1-833-387-9734.

This Patient Information has been approved by the U.S. Food and Drug Administration. Approved: 8/2020

Reference ID: 4653194

